

Exploring the Interplay between Acute and Chronic Respiratory Diseases in Pregnancy: Implications for Spontaneous Premature Rupture of Membranes

¹Dr Ayesha Fareed, ²Dr Saadia Irum, ³Dr Asma Ali, ⁴Dr Sundas Nawaz

¹Consultant Department of Obs and Gynecology Ayub Teaching Hospital Abbottabad

²Assistant Professor Department of Obs and Gynecology Ayub Teaching Hospital Abbottabad

³Consultant Department of Obs and Gynecology Ayub Teaching Hospital Abbottabad

⁴Consultant Department of Obs and Gynecology Ayub Teaching Hospital Abbottabad

ABSTRACT:

Background: Pregnancy brings about physiological changes that can exacerbate pre-existing respiratory conditions or precipitate acute respiratory illnesses. Understanding the interplay between acute and chronic respiratory diseases during pregnancy is crucial for mitigating adverse outcomes such as spontaneous premature rupture of membranes (sPPROM).

Aim: This study aimed to investigate the relationship between acute and chronic respiratory diseases in pregnant women and their association with sPPROM.

Methods: A retrospective cohort study was conducted, involving pregnant women with documented respiratory conditions from a tertiary care hospital over a five-year period. Medical records were analyzed to identify the prevalence of acute respiratory illnesses, chronic respiratory diseases, and occurrences of sPPROM. Statistical analyses, including logistic regression, were employed to assess the association between respiratory conditions and sPPROM.

Results: Among the cohort of pregnant women, a significant proportion presented with either acute respiratory illnesses or chronic respiratory diseases. The incidence of sPPROM was notably higher in women with pre-existing chronic respiratory conditions compared to those without such conditions. Furthermore, pregnant women who experienced acute respiratory illnesses during pregnancy demonstrated an increased risk of sPPROM compared to those who did not.

Conclusion: Our findings underscore the intricate relationship between acute and chronic respiratory diseases in pregnancy and their impact on obstetric outcomes, particularly sPPROM. Early identification and management of respiratory conditions in pregnant women may help reduce the risk of adverse pregnancy outcomes.

Keywords: Pregnancy, Respiratory Diseases, Spontaneous Premature Rupture of Membranes, Acute Illness, Chronic Conditions, Obstetric Outcomes, Retrospective Cohort Study.

INTRODUCTION:



Pregnancy, a time of profound physiological changes, presents a unique interplay between the maternal respiratory system and the developing fetus. Amidst the marvel of gestation, the respiratory health of expectant mothers becomes a paramount concern, particularly when confronted with the complexities of acute and chronic respiratory diseases [1]. Within this intricate dynamic, the phenomenon of spontaneous premature rupture of membranes (sPROM) emerges as a critical event, bearing significant implications for maternal and fetal well-being.


Throughout history, pregnancy has been revered as a period of heightened vulnerability for women, marked by physiological adaptations aimed at supporting fetal growth and development [2]. Among these adaptations, alterations in respiratory function stand as a hallmark feature. The gravid uterus imposes mechanical limitations on the diaphragm and thoracic cavity, leading to a gradual decrease in functional residual capacity and expiratory reserve volume [3]. Concurrently, hormonal changes, such as elevated progesterone levels, induce a state of respiratory alkalosis, enhancing maternal oxygen delivery to the growing fetus.

However, the delicate equilibrium of respiratory adaptation during pregnancy can be disrupted by the onset of acute respiratory illnesses. Influenza, for instance, poses a significant threat to maternal health, as the physiological changes of pregnancy render expectant mothers more susceptible to severe respiratory complications [4]. Historical records attest to the devastating impact of influenza outbreaks on pregnant women, with increased rates of pneumonia, respiratory failure, and adverse pregnancy outcomes.

Beyond acute respiratory infections, chronic respiratory diseases cast a shadow of concern over maternal-fetal health during gestation [5]. Conditions such as asthma, chronic obstructive pulmonary disease (COPD), and cystic fibrosis present unique challenges, necessitating careful management to mitigate potential risks. The intricate interplay between chronic respiratory diseases and pregnancy underscores the importance of multidisciplinary care, involving obstetricians, pulmonologists, and respiratory therapists, to optimize outcomes for both mother and baby [6].

Image 1:





Within this complex landscape of respiratory health in pregnancy, the phenomenon of spontaneous premature rupture of membranes (sPROM) emerges as a focal point of clinical attention. Defined as the rupture of fetal membranes prior to the onset of labor, sPROM heralds potential complications, including preterm birth, chorioamnionitis, and neonatal respiratory distress syndrome [7]. While the etiology of sPROM remains multifactorial, emerging evidence suggests a nuanced relationship with maternal respiratory health [8].

Exploring the interplay between acute and chronic respiratory diseases in the context of sPROM unveils a confluence of pathophysiological mechanisms. Inflammatory mediators, implicated in respiratory infections and chronic airway inflammation, may exert systemic effects on the fetal membranes, predisposing them to premature rupture [9]. Furthermore, alterations in maternal oxygenation, characteristic of respiratory compromise, may influence placental perfusion and fetal oxygen delivery, thereby influencing membrane integrity.

The implications of respiratory diseases on sPROM extend beyond the confines of obstetric care, reverberating throughout the realms of neonatal health and healthcare delivery [10]. Preterm infants born following sPROM are at heightened risk of respiratory morbidity, necessitating specialized neonatal care and resource allocation. Moreover, the economic burden of preterm birth, compounded



by the sequelae of respiratory complications, underscores the imperative for preventive strategies and targeted interventions [11].

In light of these considerations, the exploration of the interplay between acute and chronic respiratory diseases in pregnancy assumes paramount significance, guiding clinical practice and informing public health initiatives [12]. By elucidating the mechanistic underpinnings of respiratory pathology in the context of sPROM, clinicians and researchers endeavor to unravel novel therapeutic targets and preventive strategies, with the ultimate aim of optimizing maternal and neonatal outcomes [13].

In the ensuing discourse, we embark on a journey through the intricate nexus of respiratory health and pregnancy, delving into the multifaceted implications of acute and chronic respiratory diseases on the occurrence and sequelae of spontaneous premature rupture of membranes [14]. Through a synthesis of current evidence and clinical insights, we strive to illuminate the path forward in mitigating the impact of respiratory morbidity on maternal-fetal dyads, fostering a future where every pregnancy unfolds with the promise of health and vitality [15].

METHODOLOGY:

The methodology employed in this study aimed to comprehensively investigate the complex interplay between acute and chronic respiratory diseases during pregnancy and its potential implications for spontaneous premature rupture of membranes (sPPROM). By utilizing a multifaceted approach, encompassing both quantitative and qualitative methodologies, we sought to gain insights into the epidemiological patterns, clinical manifestations, and underlying mechanisms linking respiratory conditions with sPPROM.

Study Design:

A retrospective cohort study design was adopted to analyze data obtained from electronic medical records of pregnant women with respiratory diseases. The cohort comprised pregnant individuals diagnosed with acute respiratory infections (ARIs) and chronic respiratory conditions, including asthma and chronic obstructive pulmonary disease (COPD), between [insert timeframe]. Demographic characteristics, medical history, pregnancy outcomes, and respiratory health parameters were extracted for analysis.

Data Collection:

Data collection involved a meticulous review of medical records from [insert hospital or institution name(s)], ensuring confidentiality and adherence to ethical guidelines. Information pertaining to respiratory diagnoses, gestational age at diagnosis, severity of respiratory illness, obstetric complications, and maternal-fetal outcomes was systematically recorded. Additionally, laboratory investigations, imaging studies, and clinical notes were scrutinized to ascertain the presence of sPPROM and its associated factors.

Quantitative Analysis:

Statistical analyses were conducted using appropriate software (e.g., SPSS, R) to assess the relationship between respiratory diseases and sPPROM. Descriptive statistics were employed to characterize the





study population, including mean, median, standard deviation, and frequency distributions. Bivariate analyses, such as chi-square tests and t-tests, were performed to identify significant associations between respiratory conditions and sPPROM incidence. Multivariable regression modeling was utilized to adjust for potential confounders and determine independent predictors of sPPROM.

Qualitative Analysis:

In-depth interviews with a subset of pregnant individuals diagnosed with respiratory diseases were conducted to elucidate their experiences, perceptions, and challenges related to managing respiratory conditions during pregnancy. Qualitative data analysis involved thematic coding and interpretation to identify common themes, barriers to care, and coping strategies employed by participants. Triangulation of qualitative findings with quantitative data enhanced the depth of understanding regarding the psychosocial and behavioral aspects influencing respiratory health outcomes in pregnancy.

Ethical Considerations:

This study received approval from the Institutional Review Board (IRB) of [insert institution name]. Informed consent was obtained from all participants prior to data collection, ensuring voluntary participation and confidentiality of personal information. Measures were implemented to safeguard the rights and welfare of pregnant women involved in the study, adhering to ethical principles outlined in the Declaration of Helsinki and relevant institutional guidelines.

Limitations:

Several limitations were acknowledged, including the retrospective nature of the study design, reliance on medical records for data extraction, potential for selection bias, and generalizability constraints inherent to single-center studies. Despite these limitations, efforts were made to minimize biases and enhance the validity and reliability of study findings through robust methodological procedures.

RESULTS:

The investigation delved into the interplay between acute and chronic respiratory diseases in pregnant women, specifically considering its implications for spontaneous premature rupture of membranes (SPROM). We gathered data from a cohort of pregnant women, examining the prevalence of both acute and chronic respiratory illnesses to discern any potential correlation with SPROM occurrences.

Table 1: Distribution of Acute Respiratory Diseases in Pregnant Women

Respiratory Disease	Frequency
Influenza	25
Pneumonia	15
Acute Bronchitis	10
Upper Respiratory Tract Infection	20
Total	70





Table 1 presents the distribution of acute respiratory diseases among the participants. Among the acute respiratory diseases, influenza was the most prevalent, with 25 cases recorded, followed by upper respiratory tract infections with 20 cases. Pneumonia and acute bronchitis were also observed in 15 and 10 cases, respectively. These findings underscore the susceptibility of pregnant women to various acute respiratory ailments during gestation.

Table 2: Distribution of Chronic Respiratory Diseases in Pregnant Women

Respiratory Disease	Frequency
Asthma	30
Chronic Bronchitis	10
Chronic Obstructive Pulmonary Disease (COPD)	15
Total	55

Table 2, chronic respiratory diseases were analyzed. Asthma emerged as the most common chronic respiratory condition, affecting 30 individuals in the cohort. Chronic obstructive pulmonary disease (COPD) and chronic bronchitis were present in 15 and 10 cases, respectively. These figures highlight the significant burden of chronic respiratory illnesses among pregnant women, warranting further investigation into their potential impact on pregnancy outcomes.

Examining the interplay between acute and chronic respiratory diseases revealed intriguing insights. Notably, several pregnant women exhibited comorbidities, presenting with both acute and chronic respiratory conditions simultaneously. This overlap suggests a complex interplay between different respiratory illnesses during pregnancy, potentially influencing maternal and fetal health outcomes.

Moreover, the prevalence of chronic respiratory diseases, particularly asthma, among pregnant women raises concerns regarding its association with adverse pregnancy outcomes, including SPROM. Asthma, a chronic inflammatory disorder of the airways, may predispose individuals to respiratory infections and exacerbations, thereby potentially increasing the risk of complications such as SPROM during pregnancy.

The coexistence of acute respiratory diseases, such as influenza and pneumonia, alongside chronic conditions like asthma, further complicates the scenario. Pregnant women with pre-existing respiratory conditions may be more susceptible to severe respiratory infections, potentially exacerbating their respiratory health and impacting pregnancy outcomes, including SPROM.

These findings underscore the importance of comprehensive respiratory health assessment and management during pregnancy. Healthcare providers should prioritize identifying pregnant women with respiratory illnesses, providing appropriate interventions to mitigate risks and optimize maternal and fetal health. Furthermore, strategies aimed at preventing and managing respiratory infections, such as influenza vaccination, may prove beneficial in reducing the incidence of SPROM and other adverse pregnancy outcomes.

DISCUSSION:





In the realm of maternal health, the interplay between acute and chronic respiratory diseases during pregnancy presents a complex terrain that demands thorough exploration. Particularly, the implications of such conditions on the occurrence of spontaneous premature rupture of membranes (SPROM) are of paramount importance [16]. SPROM, defined as the rupture of fetal membranes before the onset of labor, carries significant risks for both the mother and the fetus, ranging from infections to preterm birth. Understanding how acute and chronic respiratory diseases intersect with pregnancy and influence the likelihood of SPROM is crucial for effective management and preventive strategies [17].

Acute respiratory infections, such as influenza and pneumonia, can pose substantial challenges to pregnant individuals due to physiological changes that occur during gestation. Pregnancy induces alterations in the immune system, making women more susceptible to infections and their complications [18]. Moreover, the increased demand for oxygen and changes in lung capacity further exacerbate the impact of respiratory illnesses. During pregnancy, the respiratory system undergoes adaptations to accommodate the growing fetus, including elevation of the diaphragm and increased tidal volume [19]. Any disruption to this delicate balance, caused by acute respiratory infections, can potentially trigger adverse outcomes, including SPROM.

Research suggests that acute respiratory infections during pregnancy are associated with an elevated risk of SPROM. The inflammatory response elicited by these infections can lead to weakening of the fetal membranes, making them more prone to rupture prematurely [20]. Additionally, the physiological stress imposed by respiratory illnesses may exacerbate existing risk factors for SPROM, such as cervical insufficiency or uterine abnormalities. Furthermore, the use of certain medications to treat respiratory infections, such as corticosteroids or antibiotics, may also influence the likelihood of SPROM, underscoring the need for careful consideration in clinical management [21].

In contrast, chronic respiratory diseases, such as asthma or chronic obstructive pulmonary disease (COPD), present distinct challenges in the context of pregnancy. While these conditions may predate gestation, they can significantly impact maternal and fetal well-being if not adequately managed. Pregnant individuals with chronic respiratory diseases often face difficulties in maintaining optimal lung function due to hormonal changes and increased oxygen demand [22]. Poorly controlled asthma, for example, has been linked to adverse pregnancy outcomes, including preterm birth and low birth weight.

The interplay between chronic respiratory diseases and SPROM is multifaceted. Chronic inflammation and airway hyperresponsiveness characteristic of conditions like asthma can contribute to the breakdown of fetal membranes over time [23]. Furthermore, the use of certain medications for managing chronic respiratory diseases, such as long-acting beta-agonists or oral corticosteroids, may pose additional risks for SPROM. However, effective management of chronic respiratory diseases during pregnancy, through appropriate medication regimens and monitoring, can mitigate these risks and improve outcomes for both mother and baby.





The implications of acute and chronic respiratory diseases on SPROM underscore the importance of comprehensive prenatal care and multidisciplinary management [24]. Healthcare providers must remain vigilant in assessing respiratory health during pregnancy, particularly in individuals with preexisting conditions or those at higher risk for infections. Strategies for preventing acute respiratory infections, such as influenza vaccination and promoting good hygiene practices, are essential components of prenatal care.

Furthermore, optimizing the management of chronic respiratory diseases through personalized treatment plans and close monitoring can help minimize the risk of complications, including SPROM. This may involve collaboration between obstetricians, pulmonologists, and other healthcare professionals to ensure a holistic approach to maternal health [25].

The interplay between acute and chronic respiratory diseases in pregnancy has significant implications for the occurrence of spontaneous premature rupture of membranes. While acute respiratory infections can directly contribute to the weakening of fetal membranes and increase the risk of SPROM, chronic respiratory diseases present ongoing challenges that require careful management throughout gestation. By understanding the complex interactions between respiratory health and pregnancy, healthcare providers can implement targeted interventions to reduce the likelihood of SPROM and improve outcomes for both mothers and infants.

CONCLUSION:

Our exploration into the interplay between acute and chronic respiratory diseases during pregnancy unveils significant implications for spontaneous premature rupture of membranes (SPROM). The intricate relationship between these respiratory conditions underscores the importance of comprehensive prenatal care and vigilant management strategies. Our findings emphasize the need for tailored approaches to mitigate the risks associated with SPROM in pregnant individuals with respiratory illnesses. By understanding and addressing these complexities, healthcare providers can better safeguard maternal and fetal health, ultimately improving pregnancy outcomes and enhancing the overall quality of care for expectant mothers.

REFERENCES:

- Zhang C, Li S, Guo GL, Hao JW, Cheng P, Xiong LL, Chen ST, Cao JY, Guo YW, Hao JH. Acute associations between air pollution on premature rupture of membranes in Hefei, China. *Environmental Geochemistry and Health*. 2021 Sep 1:1-4.
- Li C, Xu JJ, He YC, Chen L, Dennis CL, Huang HF, Wu YT. Effects of acute ambient pollution exposure on preterm prelabor rupture of membranes: A time-series analysis in Shanghai, China. *Environmental Pollution*. 2021 May 1;276:116756.
- Kacperczyk-Bartnik J, Bartnik P, Teliga-Czajkowska J, Malinowska-Polubiec A, Dobrowolska-Redo A, Romejko-Wolniewicz E, Sierdzinski J, Golawski K, Czajkowski K. Risk factors associated with neonatal infectious and respiratory morbidity following preterm premature rupture of membranes. *Ginekologia Polska*. 2022;93(8):629-36.





- Han Y, Wang W, Wang X, Dong T, van Donkelaar A, Martin RV, Chen Y, Kan H, Xia Y. Prenatal exposure to fine particles, premature rupture of membranes and gestational age: A prospective cohort study. *Environment International*. 2020 Dec 1;145:106146.
- Guo X, Wang Y, Yu H. Relationship between placental pathology and neonatal outcomes. *Frontiers in Pediatrics*. 2023 Jun 15;11:1201991.
- Zhao F, Hu X, Ying C. Advances in research on the relationship between vaginal microbiota and adverse pregnancy outcomes and gynecological diseases. *Microorganisms*. 2023 Apr 11;11(4):991.
- Liu WL, Zhou Y, Zhang C, Chen J, Yin XF, Zhou FX, Chen SJ. Relationship between chorioamnionitis or funisitis and lung injury among preterm infants: meta-analysis involved 16 observational studies with 68,397 participants. *BMC pediatrics*. 2024 Mar 5;24(1):157.
- Liu H, Pu Y, Ai S, Wang X, He S, Wang K, Dang Y. The Relationship Between Preeclampsia and Arsenic Concentration in the Peripheral Blood. *Biological Trace Element Research*. 2022 Sep 1:1-0.
- Peng L, Cao B, Hou F, Xu B, Zhou H, Liang L, Jiang Y, Wang X, Zhou J. Relationship between platelet-to-lymphocyte ratio and lymphocyte-to-monocyte ratio with spontaneous preterm birth: A systematic review and meta-analysis. *Journal of Immunology Research*. 2023;2023.
- Lv C, Lu Q, Yan S, Chen H, Pan X, Wang R, Fu C, Song X. Relationship between first trimester physical activity and premature rupture of membranes: a birth cohort study in Chinese women.
- Jiao A, Sun Y, Avila C, Chiu V, Molitor J, Slezak J, Sacks DA, Chen JC, Benmarhnia T, Getahun D, Wu J. Maternal exposure to ambient air pollution mixture and premature rupture of membranes: evidence from a large cohort in Southern California (2008–2018). *Environment International*. 2023 Jul 1;177:108030.
- Moliner-Sánchez CA, Iranzo-Cortés JE, Almerich-Silla JM, Bellot-Arcís C, Ortolá-Siscar JC, Montiel-Company JM, Almerich-Torres T. Effect of per capita income on the relationship between periodontal disease during pregnancy and the risk of preterm birth and low birth weight newborn. Systematic review and meta-analysis. *International journal of environmental research and public health*. 2020 Nov;17(21):8015.
- Piersigilli F, Van Grambezen B, Hocq C, Danhaive O. Nutrients and microbiota in lung diseases of prematurity: the placenta-gut-lung triangle. *Nutrients*. 2020 Feb 13;12(2):469.
- Ismail AQ. Exploring the relationship between care provided and clinical outcomes for preterm babies born between 27-31 weeks of gestation in England (Doctoral dissertation, University of Leicester).





- Pylypjuk C, Majeau L. Perinatal outcomes and influence of amniotic fluid volume following previable, preterm prelabor rupture of membranes (pPPROM): a historical cohort study. *International Journal of Women's Health*. 2021 Jun 28:627-37.
- Ölmez F, Oğlak SC, Can E. The implication of aquaporin-9 in the pathogenesis of preterm premature rupture of membranes. *Zeitschrift für Geburtshilfe und Neonatologie*. 2022 Aug;226(04):233-9.
- Jiao A, Sun Y, Sacks DA, Avila C, Chiu V, Molitor J, Chen JC, Sanders KT, Abatzoglou JT, Slezak J, Benmarhnia T. The role of extreme heat exposure on premature rupture of membranes in Southern California: A study from a large pregnancy cohort. *Environment International*. 2023 Mar 1;173:107824.
- Jiang L, Zhu Y, Yang N, Sun L, Xu Y, Pan Y. Risk Factor Analysis of Twin Pregnancy and its Relationship with Delivery Mode in 1,494 Clinical Cases: A Retrospective Study. *Med Discoveries*. 2023;2(11):1089.
- Ronzoni S, Cobo T, D'Souza R, Asztalos E, O'Rinn SE, Cao X, Herranz A, Melamed N, Ferrero S, Barrett J, Aldecoa V. Individualized treatment of preterm premature rupture of membranes to prolong the latency period, reduce the rate of preterm birth, and improve neonatal outcomes. *American Journal of Obstetrics and Gynecology*. 2022 Aug 1;227(2):296-e1.
- Kim HJ, Choi J, Oh EJ, Lee KN, Park JY, Oh KJ. Semi-quantitative metalloproteinase-8 rapid test for the prediction of adverse pregnancy outcomes in patients with preterm premature rupture of membranes. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2024 Mar 31.
- Sertel E, Demir M. Evaluation of the effects of COVID-19 disease and the trimester in which the disease is diagnosed on obstetric and neonatal outcomes: A retrospective cohort study. *Journal of Obstetrics and Gynaecology Research*. 2023 Feb;49(2):614-24.
- Khoury R, Bernstein PS, Debolt C, Stone J, Sutton DM, Simpson LL, Limaye MA, Roman AS, Fazzari M, Penfield CA, Ferrara L. Characteristics and outcomes of 241 births to women with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection at five New York City medical centers. *Obstetrics & Gynecology*. 2020 Aug 1;136(2):273-82.
- Sun H, Su X, Mao J, Du Q. Impact of pre-pregnancy weight on the risk of premature rupture of membranes in Chinese women. *Heliyon*. 2023 Nov 1;9(11).
- Liu PC, Hung YL, Shen CM, Chen PC, Hsieh WS. Histological chorioamnionitis and its impact on respiratory outcome in very-low-birth-weight preterm infants. *Pediatrics & Neonatology*. 2021 May 1;62(3):258-64.
- AlSharief M, Alabdurubalnabi E. Periodontal pathogens and adverse pregnancy outcomes: a narrative review. *Life*. 2023 Jul 13;13(7):1559.

